

## Frequently asked questions: Influenza (flu) information for parents

### Seasonal influenza 2013-2014

Version 1.1

9<sup>th</sup> December 2013

*The following advice is for parents of children in all educational institutions, including crèches, childcare, schools, and third level institutions. Unless otherwise mentioned 'educational institution' applies to all of the above.*

We are now experiencing our annual flu season and influenza A and B are circulating in the community.

#### ***Is influenza dangerous?***

Most people infected with the influenza virus have a mild to moderate illness, but some have more severe illness.

#### ***What are the symptoms of influenza?***

The symptoms of flu include:

- Temperature over 38 °C/100 °F that begins suddenly **and** some of the following:
  - Dry cough
  - Sore throat
  - Muscle aches and pains
  - Headache
  - Runny nose
  - Severe weakness and fatigue
  - Vomiting/diarrhoea (in some cases)

Symptoms	Influenza	Common Cold
Onset	Sudden	Slow
Fever	Characteristically High ( $\geq 38^{\circ}\text{C}$ or $100^{\circ}\text{F}$ )	Rare
Headache	Prominent	Rare
General aches and pains	Usual, often severe	Rare
Fatigue, weakness	Can be prolonged for a number of weeks	Quite mild
Extreme exhaustion	Early and prominent	Never
Runny nose	Common	Common
Sneezing	Common	Usual
Sore throat	Common	Common
Cough	Common, can be severe	Mild to moderate, hacking cough
Diarrhoea, vomiting	Sometimes	Not associated with the common cold in adults

### ***How does influenza spread?***

Flu virus spreads from person to person mainly through the coughing or sneezing of a sick person. Flu virus may also be spread when a person touches something that is contaminated with the virus (for example a tissue or door handle touched by the infected person) and then touches his or her eyes, nose or mouth.

### ***What should I do as a parent?***

#### **Two important actions to protect your family**

1. Be aware of the symptoms of flu-like illness and know where to seek medical care. See below.

- Teach your children to avoid touching their eyes, nose and mouth.
- Teach your children to stay at least 1 metre/3 feet away from people who are sick.
- Children who are sick should always stay home from the school. If they have influenza they should stay away from the educational institution for 5 days from the onset of symptoms.
- Wash hard surfaces such as kitchen worktops, door handles, etc with a normal household cleaner as the virus can live on these surfaces. Do this frequently.

***What should I do if my child gets sick?***

If your child gets sick with a flu-like illness as described above you should:

- Keep your child at home and away from others as much as is possible to avoid spreading infection to others. If they are sick with flu they should stay home for 5 days from the onset of symptoms.
- Give your child simple anti-fever medication such as paracetamol or ibuprofen (NB: aspirin should NOT be given to children under 16 years of age) and ensure that they drink plenty of fluids.
- If you think your child needs to see the GP because they have severe symptoms, remember to ring your GP first.
- If your child is in a high risk group for complications of flu contact your GP, even if their symptoms are mild (High risk groups include people with: chronic lung, heart, kidney, liver, or neurological disease; immunosuppression ( i.e. weakened immune system whether caused by disease or treatment); diabetes mellitus; people aged 65 years and older; children less than 2 years of age; children with any condition (e.g. spinal cord injury, seizure disorder or other neuromuscular disorder) that can compromise lung function, especially those attending special schools/day centres; people on medication for asthma, severely obese people (BMI  $\geq 40$ ), pregnant women and people with haemoglobinopathies)
- Teach your child good health habits, as above.
- Ensure that all household surfaces that are touched by hands are kept clean, especially bedside tables, surfaces in bathrooms and kitchens and children's toys. Such surfaces should be wiped regularly with a household disinfectant according to directions on the product label.

- Children should wash their hands frequently with soap and water. Where soap and water is not readily available alcohol based hand gel can be used.
- Children should avoid touching their eyes, nose and mouth.
- Wash hard surfaces such as kitchen worktops, door handles, etc frequently with a normal household cleaner as the virus can live on these.

Educational institutions should put up posters on respiratory etiquette and hand hygiene and ensure that children have access to suitable hand washing facilities. Posters can be found on the HPSC website as indicated below.

### ***Is there a vaccine against seasonal influenza?***

A vaccine is an injection which prevents a person getting a particular disease. It works by strengthening the body's immune system. The vaccine must be given before the person is infected with the disease.

Vaccination is recommended for people at high risk of complications.

### ***Who are the high risk groups that should receive seasonal influenza vaccine?***

1. Those aged 65 years or older.
2. Adults and children over 6 months of age with chronic illness requiring regular medical follow-up e.g. chronic heart disease, chronic liver disease, chronic neurological disease (including multiple sclerosis and hereditary and degenerative disorders of the central nervous system), chronic renal failure chronic respiratory disease ( including cystic fibrosis, moderate or severe asthma, and bronchopulmonary dysplasia), diabetes mellitus, haemoglobinopathies
3. Those with immunosuppression (weakened immune system) due to disease or treatment, including asplenia (no spleen) or splenic dysfunction.
4. Those with any condition that can compromise lung function (e.g. spinal cord injury, seizure disorder, or other neuromuscular disorder) especially those attending special schools/day centres.
5. Those with Down syndrome.
6. Children with moderate to severe neurodevelopmental disorders such as cerebral palsy and intellectual disability.

11. All pregnant women at any stage of pregnancy.

12. Those who have close, regular contact with pigs, poultry or water fowl.

***Are there medicines to treat influenza?***

Yes, there are medicines known as anti-virals that can be used to treat flu. However, as most cases of influenza will be mild, anti-viral treatment will only be necessary in a small proportion of cases. Doctors will assess each case but the following groups are the ones most likely to require treatment with anti-virals:

- Patients who appear to have severe symptoms or
- Patients who are in defined high risk groups as above.

***Further information***

Posters:

<http://www.hpsc.ie/hpsc/A-Z/Gastroenteric/Handwashing/PostersVideos/>

<http://www.hpsc.ie/hpsc/A-Z/Respiratory/Influenza/SeasonalInfluenza/Infectioncontroladvice/>

Guidance on seasonal influenza:

<http://www.hpsc.ie/hpsc/A-Z/Respiratory/Influenza/SeasonalInfluenza/>